

Titan arum (*Amorphophallus titanum*) flowers in New Zealand for the first time

Ewen K. Cameron¹

On the 30 November 2013 a titan arum (*Amorphophallus titanum*) flowered in the Tropical House at the Wintergarden in the Auckland Domain, attracting large crowds (Fig. 1) as it was the first time this species has flowered in New Zealand. It belongs to the arum family (Araceae) and the 'flower' as such is not a single flower, but a cluster of flowers – an inflorescence. The central fleshy spike, the spadix, has a zone of male flowers and below them a separate zone of female flowers; the spadix is wrapped in a leaf-like bract, the spathe, which is open above allowing pollination (Fig. 2). Well-known cultivated members of the family include philodendrons (*Philodendron*) and calla or arum lilies (*Zantedeschia*). The name titan arum was coined by David Attenborough during the filming of the *Private Life of Plants* series (1995). Titan arum is often considered the largest inflorescence of herbaceous plants (cf. the palm genus *Corypha*) but *Amorphophallus gigas* is alleged to be taller (Mabberley, 2008).



Fig. 1 The local radio kept announcing the exciting news and c. 10,000 visitors flocked to the Wintergarden in the Auckland Domain to see the first titan arum flowering in New Zealand on Sunday 1 December 2013. The queue double-looped around the courtyard between the two glasshouses. All photos by EKC.

The titan arum was discovered and described in a different genus (*Conophallus*) in 1878 by the Italian botanist Odoardo Beccari and

transferred to *Amorphophallus* the following year. There are about 150 species in the genus *Amorphophallus* occurring in the Old World tropics (Mabberley, 2008). A translation of the scientific name:

amorphous (Greek) – shapeless or deformed

phallus – penis (referring to the spadix)

titanus – very large.




Fig. 2 A stylised diagram of an arum inflorescence (image improved by Joshua Salter).

Titan arum is native to the tropical rainforests of Sumatra (Indonesia) and is locally known as *bunga bangkai*, roughly translated as 'corpse flower'. It is classed as 'Vulnerable' on the IUCN Red List of threatened plants – much of its natural habitat is being cleared for oil palm plantations. The underground tuber, a corm, usually weighs up to 50 kg, but one at the Botanic Garden at Bonn (Germany) weighed 117 kg (Liptrot, 2013). Each year the plant usually grows one leaf that can be up to 6 m tall (Fig. 3) and lasts for over a year. It has been suggested that the lichen-like patches on the petiole (Fig. 4) are mimicry to prevent collision by animals² which could easily damage the trunk-like petioles (Hejnowicz and Barthlott, 2005). The corm enters a short

dormant period (unusual in rainforest species) before producing another leaf, or every few years it flowers instead of producing a leaf.

"The flowering structure grows 10 cm per day [and may reach 3 m tall], then heats up to human body temperature to broadcast its smell like smoke from a chimney. A Kew chemist found that its signature scent of "rotting animal in a pan of sauerkraut" is due to dimethyl sulphides. The stench attracts insects [carrion beetles and flies] to carry pollen between titan arum plants" (a label at Kew Gardens, June 2011).



Fig. 3 A single giant, umbrella-like compound leaf of titan arum. Princess of Wales Conservatory at Kew Gardens, England, 13 June 2011.



Fig. 4 Petiole (18 cm diam.) of a 4 m tall leaf. Photo: 7 April 2014. Figs. 4–10 taken in the Tropical House (TH), Wintergarden, Auckland Domain.

¹ Curator Botany, Auckland War Memorial Museum, The Domain, Private Bag 92018, Victoria Street West, Auckland 1142, New Zealand; ecameron@aucklandmuseum.com

² The reasoning being that the lichen appearance makes the petiole look like a woody trunk and therefore large animals avoid hitting them.



Fig. 5 Titan arum inflorescence 2.5 m tall – the spathe started opening 3 hrs after photo taken. Photo: 11 am, 30 November 2013.



Fig. 6 Fully open titan arum, scent in weak pulses; the 2-hour wait was worth it. Photo: 8 pm, 1 December 2013.



Fig. 7 Spathe closed, no scent. Photo: 8.30 am, 3 December 2013.



Fig. 8 Spadix collapsed over previous night. Photo: 2 pm, 4 December 2013.



Fig. 9 Spadix withered and top of spathe wilted. Photo: 8 December 2013.



Fig. 10 New leaf (2.4 m tall) of plant that flowered on 30 November 2013. Photo: 7 April 2014 with Eveline Perl, gardener at TH.

Table 1 The diary of the Auckland Domain titan arum.

2008	One of two corms c. 2 years old acquired in a plant exchange by Domain Nursery.
11 Nov 2013	A bud emerged from the 50 kg corm after 6 months dormancy.
18 Nov 2013	Top of white spathe visible, (therefore not a leaf).
30 Nov 2013	2 pm the spathe began to open in the Tropical House maintained at 28°C (Fig. 5).
1 Dec 2013	Spathe fully open (Fig. 6) and smelly in pulses; the Wintergarden gates opened at 6.30 am, closed at 7.30 pm with the last visitors leaving at 9.15 pm; the day's total c. 10,000 visitors.
2 Dec 2013	Spathe still open, but wilting a little, scent very faint.
3 Dec 2013	Spathe closed by 8.30 am, no scent (Fig. 7).
4 Dec 2013	Spadix collapsed overnight (Fig. 8).
8 Dec 2013	Spadix withered (Fig. 9).
20 Dec 2013	Spathe now withered.
3 Feb 2014	Totally dried up, plant retained in Tropical House.
10 Feb 2014	A new leaf shoot emerged.
7 Apr 2014	New leaf only c. 2.4 m tall and yet to unfold (Fig. 10).

The female flowers are receptive first, the male flowers releasing pollen the next day; in nature, this timing ensures cross-pollination with another flowering plant. However, solitary cultivated blooms occasionally manage to self-pollinate. If flowers are successfully pollinated, the surrounding spathe eventually falls off, exposing the maturing seeds; when ripe, the cherry-sized fruits turn a bright orange-red, a colour attractive to birds (including hornbills) which in nature pick the berries off and dispersed the seed (http://botit.botany.wisc.edu/Titan_Arum_Archive/).

The first plant in the world to flower in cultivation was at the Royal Botanic Gardens Kew in 1889 to a riotous reception (from seed sent by Beccari). The second was in 1894 at Land's Plantentuin Buitenzorg, Kebun Raya Bogor in Java (Indonesia). Up until 1993 there had only been 22 publicised flowering events from cultivated plants worldwide, but since then there has been another 143 publicised blooms, with the Auckland one being the most recent (http://en.wikipedia.org/wiki/List_of_publicised_titan_arum_blooms_in_cultivation).

The first one to flower in Australia was in Cairns (Fletcher Botanic Gardens) in 2003, then at the Royal Botanic Gardens Sydney the following year, and more recently at the Royal Botanic Gardens Melbourne in December 2012. There are plenty of time lapse sequences of the inflorescence opening online (check *YouTube*).

The flowering of the one at the Auckland Domain Wintergarden was well-advertised and the public came flocking in. On Saturday morning (Fig. 5) the staff suspected it wouldn't open until the next day.

On Sunday I had the Auckland Botanical Society picnic to attend at Tawharanui Regional Park. After the picnic and bush walk a few of us decided to check the titan arum out. We arrived c. 6 pm, and parking in the Domain was at a premium. We patiently queued with everyone else for nearly two hours (Fig. 1) to be rewarded with the colourful spathe magnificently open (Fig. 6) and pulses of quite manageable sulphide-like aroma. It was wonderful to participate in what I suspect was the largest one-day botanical event in the history New Zealand. Over the next few weeks the plant withered up (Figs. 7–9, Table 1); and by the end of January there appeared to be no living tissue left above the ground. After a very short dormancy, surprisingly then, on the 10 February a new leaf shoot emerged from the centre of the corm. It has since developed quite slowly and on 7 April it was only c. 2.4 m tall, with a petiole diameter 10 cm at 1 m, and was yet to unfold (Fig. 10). Compare this with another titan arum plant in the same glasshouse, whose new leaf appeared at the same time as the one on the plant that flowered, and on 7 April was 4 m tall, with a petiole diameter 18 cm at 1 m and the leaf blade expanded (similar to the one in Fig. 3). Presumably the flowering of the other plant has robbed its corm of its reserves so that it must

now build them up. However, the prospects look good for a flowering next year of the second plant – don't miss it if it does.

Acknowledgements

I thank the Auckland Domain Nursery and Wintergarden staff: Eveline Perl, Paul Novotny and Melanie James for information about the titan arums under their care; and Joshua Salter for improving Figure 2.

References

- Hejnowicz, Z. and Barthlott, W. (2005). Structural and mechanical peculiarities of the petioles of giant leaves of *Amorphophallus* (Araceae). *American Journal of Botany* 99: 391–403.
- Liptrot, M. (2013). Corpse flower a kiwi first. *New Zealand Herald*, 1 December 2013.
- Mabberley, D.J. (2008). Mabberley's plant-book. A portable dictionary of plants, their classification and uses. Third ed. Cambridge University Press, Cambridge. 1021 p.
- Websites**
http://botit.botany.wisc.edu/Titan_Arum_Archive/ – accessed 2 November 2014.
http://en.wikipedia.org/wiki/List_of_publicised_titan_arum_blooms_in_cultivation – accessed 2 November 2014.

Reproduced with permission from the *Auckland Botanical Society Journal*, Vol. 69, No. 1, June 2014, pp. 85–88.